



## Measurement Reasonable Value - Time (metric)

<p><b>1</b> What is the most reasonable value for the given measurement?</p> <p>A</p> <p>6 d</p>	<p>The time to drive across North America</p> <p>B</p> <p>6 hr</p>	<p><b>2</b> What is the most reasonable value for the given measurement?</p> <p>The time for a decent runner to run the length of a football field</p> <table border="1"> <tbody> <tr> <td data-bbox="1169 520 1357 722">A</td> <td data-bbox="1357 520 1547 722">B</td> </tr> <tr> <td>25 s</td> <td>25 min</td> </tr> </tbody> </table>	A	B	25 s	25 min
A	B					
25 s	25 min					
<p><b>3</b> What is the most reasonable value for the given measurement?</p> <p>A</p> <p>90 min</p>	<p>The time it takes to eat dinner at a restaurant</p> <p>B</p> <p>90 s</p>	<p><b>4</b> What is the most reasonable value for the given measurement?</p> <p>A</p> <p>90 s</p> <p>B</p> <p>90 min</p>				
<p><b>5</b> What is the most reasonable value for the given measurement?</p> <p>A</p> <p>45 hr</p>	<p>The time for a subway ride across a city</p> <p>B</p> <p>45 min</p>	<p><b>6</b> What is the most reasonable value for the given measurement?</p> <p>A</p> <p>500 ms</p> <p>B</p> <p>500 s</p>				
<p><b>7</b> What is the most reasonable value for the given measurement?</p> <p>A</p> <p>200 ms</p>	<p>The time for one flap of a hummingbird's wing</p> <p>B</p> <p>200 s</p>	<p><b>8</b> What is the most reasonable value for the given measurement?</p> <p>A</p> <p>2 min</p> <p>B</p> <p>2 hr</p>				