

mobius

Pythagorean Theorem - Either Missing Length (Decimal)



1	Find the length of the missing side as a decimal value based on the Pythagorean theorem	A m=11.45	B m=8.59	Find the length of the missing side as a decimal value based on the Pythagorean theorem	A r=5.71	r=15
	10 3 m	C m=9.54	D m=10.54		C r=6.04	D r=6.71
		E m=8.54	F m=5.54	r	e r=54	F r=8.72
3	Find the length of the missing side as a decimal value based on the Pythagorean theorem	A x=7.28	B x=4.76	Find the length of the missing side as a decimal value based on the Pythagorean theorem	A r=5.31	B r=1.11
		C x=8.12	D x=8.96		C r=3.63	D r=8
		E x=6.44	F x=9		E r=1.95	F r=4.47
5	Find the length of the missing side as a decimal value based on the Pythagorean theorem	s a sed on r=7.06 r=13 missing decimal value in the Py	Find the length of the missing side as a decimal value based on the Pythagorean theorem	А m=13	в m=42	
	9 4 r	C r=8.06	D r=11.29	7 m	C m=8.38	D m=12.58
		E r=9.67	F r=5.64		E m=3.61	F m=9.22
7	Find the length of the missing side as a decimal value based on the Pythagorean theorem	A z=2.03	B z=5.39	Find the length of the missing side as a decimal value based on the Pythagorean	A z=4.51	в z=1
		C z=4.58	D z=4.55		C z=1.15	D z=6.19
		E z=6.23	r z=10	z	E z=2.83	F z=4