

mobius

Pythagorean Theorem - Either Missing Length (Decimal)



1	Find the length of the missing side as a decimal value based on the Pythagorean theorem	n=90	B n=12.61	2	Find the length of the missing side as a decimal value based on the Pythagorean theorem	A n=13.21	B n=12.37
		C n=15.13	n=19			C n=9.85	D n=10.69
		n=13.45	n=10.93			n=13	n=11.53
7	Find the length of the missing side as a decimal value based on the Pythagorean theorem	A r=9.69	r=70	4	Find the length of the missing side as a decimal value based on the Pythagorean theorem	A p=1.63	B p=6.63
		C r=12.21	D r=15.57			C p=4.64	D p=11.58
		r=8.85	r=8.01			p=22	p=5.63
5	Find the length of the missing side as a decimal value based on the Pythagorean theorem	A m=14	B m=11.61	6	Find the length of the missing side as a decimal value based on the Bythagorean theorem	A r=15.89	B r=14.21
		C m=10.77	D m=9.17			C r=13.37	D r=15.05
		E m=8.25	F m=6.57			E r=10.85	F r=16.73
7	Find the length of the missing side as a decimal value based on the Pythagorean theorem	A z=15.56	B z=17.24	6	Find the length of the missing side as a decimal value based on the Pythagorean theorem	A p=15.18	p=12.65
		C z=13.88	D z=16.4			C p=7.59	D p=16.44
		E z=22	r z=121			E p=15.65	F p=6.32