




Pythagorean Triples (Scaled) - Length of Side




1 Find the length of the missing side as a decimal value based on the Pythagorean theorem




A	B	C
1,800	2,000	1,700
D	E	F
1,600	2,100	1,000

2 Find the length of the missing side as a decimal value based on the Pythagorean theorem




A	B	C
100	130	110
D	E	F
120	140	180

3 Find the length of the missing side as a decimal value based on the Pythagorean theorem




A	B	C
200	240	170
D	E	F
250	210	400

4 Find the length of the missing side as a decimal value based on the Pythagorean theorem




A	B	C
320	140	150
D	E	F
160	220	110

5 Find the length of the missing side as a decimal value based on the Pythagorean theorem




A	B
18,000	20,000
C	D
16,000	15,000
E	F
14,000	13,000

6 Find the length of the missing side as a decimal value based on the Pythagorean theorem




A	B
10,000	15,000
C	D
16,000	12,000
E	F
18,000	9,000

7 Find the length of the missing side as a decimal value based on the Pythagorean theorem



A	B
7,000	14,000
C	D
15,000	135,000
E	F
12,000	11,000

8 Find the length of the missing side as a decimal value based on the Pythagorean theorem



A	B
10,000	7,000
C	D
12,000	8,000
E	F
65,000	13,000